

# Useful Probabilistic Inequalities

DS-563 / CD-543 @ Boston University

Spring 2023 (last updated: 2/10/2023)

## Union Bound

For any probabilistic events  $\mathcal{E}_1, \dots, \mathcal{E}_k$ ,

$$\Pr(\text{at least one of events } \mathcal{E}_1, \dots, \mathcal{E}_k \text{ has occurred}) \leq \sum_{i=1}^k \Pr(\mathcal{E}_i),$$

where  $\Pr(\mathcal{E}_i)$  denotes the probability of event  $\mathcal{E}_i$ .

In this class, we routinely use the union bound to show that we can avoid a set of bad events with good probability. For instance, consider bad events  $\mathcal{E}_1$ ,  $\mathcal{E}_2$ , and  $\mathcal{E}_3$  that can break our algorithm and occur with probability at most  $\delta/4$ ,  $\delta/5$ , and  $\delta/2$ , respectively. Then the union bound allows us to say that our algorithm works correctly with probability at least  $1 - (\delta/4 + \delta/5 + \delta/2) \geq 1 - \delta$ .

## Markov's Inequality

Let  $X$  be a non-negative random variable with  $E[X] < \infty$ . For any  $a > 0$ ,

$$\Pr(X \geq a) \leq \frac{E[X]}{a}.$$

Suppose that a generous stranger leaves an envelope with money in your mailbox every day. If on *average* there is \$100 in the envelope, how often is there at least \$200? Clearly, you cannot find this much in the envelope every day, because then the average would be at least \$200. Can you find this much 51% of the days? Again, the answer is no, because that would imply that the average would be at least  $\frac{51}{100} \cdot \$200 = 102$ , even if you assume that you get nothing on the remaining 49% of days. Markov's inequality generalizes this type of thinking to give a bound on the probability of a random variable being greater than a specific value.

Exercise: Why is the assumption that the variable is non-negative important in the above reasoning? Would it still hold if the "generous" stranger could take money from you?

## Chebyshev's Inequality

Let  $X$  be a random variable with finite expectation and variance. For any  $a > 0$ ,

$$\Pr(|X - E[X]| \geq a\sqrt{\text{Var}[X]}) \leq \frac{1}{a^2}.$$

The variance of  $X$ , i.e.,  $\text{Var}[X] = E[(X - E[X])^2]$ , is a measure how much on average  $X$  diverges from its expectation. If we have a bound on the variance of  $X$ , we can bound the probability that  $X$  significantly diverges from its expectation. This bound is very useful when  $X$  is a sum of other random variables—e.g.,  $X = \sum_{i=1}^n X_i$ —that are not fully independent. The standard proof of Chebyshev's inequality is a relatively easy application of Markov's inequality, which uses the fact that  $(X - E[X])^2$  is a non-negative variable.

## Chernoff Bound (Multiplicative Concentration)

Let  $X_1, \dots, X_n$  be independent random variables taking on values in  $[0, 1]$ . Let  $X = \sum_{i=1}^n X_i$  and let  $\mu = E[X]$ .

For any  $\delta \in [0, 1]$ ,

$$\Pr(X \leq (1 - \delta)\mu) \leq e^{-\delta^2\mu/2},$$

and

$$\Pr(X \geq (1 + \delta)\mu) \leq e^{-\delta^2\mu/3}.$$

For any  $\delta \geq 1$ ,

$$\Pr(X \geq (1 + \delta)\mu) \leq e^{-\delta\mu/3}.$$

Consider tossing an unbiased coin. Intuitively, you expect that the fraction of both heads and tails will converge to  $1/2$  as the number of trials increases. But how fast is it going to happen? This is where the Chernoff bound becomes very useful. As opposed to Chebyshev's inequality, it assumes that the variables are fully independent. This inequality can also be proved via Markov's inequality but the proof is more sophisticated.

Exercise: In the example above, what is the probability that the fraction of heads is at most  $2/5$  or at least  $3/5$  as a function of  $n$ , the number of coin tosses? Set  $X_i = 1$  if in the  $i$ -th trial the coin comes up heads, and set  $X_i = 0$ , otherwise.

## Collisions (the Birthday Paradox)

We say that there is a *collision* in a set of samples if two of them are identical.

Consider  $k$  independent samples  $x_1, x_2, \dots, x_k$  from the uniform distribution on  $\{1, \dots, n\}$ . If  $k \geq 2\lceil\sqrt{n}\rceil$ , then the probability of a collision in this set of samples is at least  $1/2$ .

Why? Suppose that there is no collision in the set of the first  $\lceil\sqrt{n}\rceil$  samples, i.e.,  $x_1, \dots, x_{\lceil\sqrt{n}\rceil}$ . Then the probability of any other sample colliding with one of them is at least  $\lceil\sqrt{n}\rceil/n \geq 1/\sqrt{n}$ . Since the samples are independent, the probability that none of the other  $\lceil\sqrt{n}\rceil$  samples collide with them is at most

$$\left(1 - \frac{1}{\sqrt{n}}\right)^{\lceil\sqrt{n}\rceil} \leq e^{-\frac{1}{\sqrt{n}} \cdot \sqrt{n}} = e^{-1} < 1/2.$$

Note 1: It can be showed that the uniform distribution minimizes the probability of a collision, so this bound holds for any distribution, not just the uniform distribution.

Note 2: This problem is referred to as the *birthday paradox*. If one performs the exact computation then a set of 23 people suffices to find a pair with the same birthday with probability more than  $1/2$ . This may seem counterintuitive, because that's much less than 365, the number of days in a typical year.

Consider  $k$  independent samples  $x_1, x_2, \dots, x_k$  from the uniform distribution on  $\{1, \dots, n\}$ . For any  $p \in [0, 1]$ , if  $k < \sqrt{2np}$ , the probability of seeing a collision is less than  $p$ .

Why? For each pair  $x_i$  and  $x_j$ , the probability that they are identical, i.e., collide, is  $\frac{1}{n}$ . Hence the expected number of identical pairs of samples is  $\binom{k}{2} \cdot \frac{1}{n} < \frac{k^2}{2n}$ . By Markov's inequality, the probability that at least one pair of samples is identical, which is equivalent to having a collision in the set of samples, is at most  $\frac{k^2}{2n}$ . If  $k < \sqrt{2np}$ , this is less than  $p$ .

In particular, this implies that if we want to see a pair of identical elements drawn with constant probability, we need  $\Omega(\sqrt{n})$  samples, i.e., the asymptotic behavior of the previous bound is tight.

## Hoeffding's Inequality

Let  $X_1, \dots, X_n$  be independent random variables such that each  $X_i \in [a_i, b_i]$ . For any  $t \geq 0$ ,

$$\Pr \left( \left| \sum_{i=1}^n (X_i - E[X_i]) \right| \geq t \right) \leq 2 \exp \left( - \frac{2t^2}{\sum_{i=1}^n (b_i - a_i)^2} \right).$$

The scenario in which this inequality is most useful in this course is the case of  $X_i$ 's being indicator variables, or more generally,  $X_i \in [0, 1]$  for all  $i \in [n]$ . In this case, the inequality becomes

$$\Pr \left( \left| \sum_{i=1}^n (X_i - E[X_i]) \right| \geq t \right) \leq 2 \exp(-2t^2/n)$$

for any  $t \geq 0$ . Alternately, we can write it as

$$\Pr \left( \left| \sum_{i=1}^n (X_i - E[X_i]) \right| \geq \epsilon n \right) \leq 2 \exp(-2\epsilon^2 n)$$

for any  $\epsilon \geq 0$ . This should look very familiar to the Chernoff bound, and in fact, one can prove a slightly weaker version of this inequality via a direct application of the Chernoff bound. The additive bound in Hoeffding's inequality is sometimes more convenient than the multiplicative bound in the Chernoff bound.

## Bonus: Non-probabilistic Inequalities

For any  $x \in \mathbb{R}$ ,  $1 + x \leq e^x$ .

## Bonus: Notation

**The set of the first  $n$  positive integers:** For any  $n \in \mathbb{N}$ ,  $[n] \stackrel{\text{def}}{=} \{1, 2, \dots, n\}$ .

**Rounding a real number up:** For any  $x \in \mathbb{R}$ ,  $\lceil x \rceil \stackrel{\text{def}}{=} \min \{y \in \mathbb{Z} : y \geq x\}$ .

**Rounding a real number down:** For any  $x \in \mathbb{R}$ ,  $\lfloor x \rfloor \stackrel{\text{def}}{=} \max \{y \in \mathbb{Z} : y \leq x\}$ .